

Download Plant Poisonings Mycotoxicoses Of Livestock In South Africa

ANIPEDIA is an online quality and invaluable scientific animal health knowledge resource providing updated information on Infectious and parasitic diseases, Plant poisonings and mycotoxicoses for veterinary, paraveterinary and allied animal and human health professionals and students.

Cestrum Laevigatum: Description
It is a much-branched shrub or tree, growing 6-15 m high in coastal regions, but inland, in areas like the Free State and the Mpumalanga provinces of South Africa, it usually attains a height of 1-2 m.

Cardiac glycoside toxicity is caused by consumption of cardiac glycoside-containing plants. These include 11 different plant families which include Apocynaceae, Asclepiadaceae, Celastraceae, Brassicaceae, Lilaceae, Moraceae, Fabaceae, Ranunculaceae, Scrophulariaceae, Sterculiaceae, and Tiliaceae. Within the 11 plant families, there are a handful of plants that are most likely to be the cause of ...

Introduction. Pneumonic pasteurellosis by definition refers to infection of the lung(s) with organisms of the genus *Pasteurella*.¹⁹ The term pneumonic pasteurellosis has served conveniently since the 1960s^{17, 138} as a means of speaking collectively of pneumonias caused by *Pasteurella haemolytica* and *Pasteurella multocida* without committing oneself as to the relative importance of the two agents.